



RECEIVED

JUL 19 2001

TECH CENTER 1600/2900

The plant of claim 16, wherein at least one of said flowers comprises a plurality of white wings.--

--19. The plant of claim 16, wherein at least one of said flowers comprises a white keel.--

--20. The plant of claim 16, wherein said plant comprises a plurality of pods whose positions on said plant are scattered.--

--21. The plant of claim 16, wherein said plant comprises a plurality of flowers and pods, said stalk, leaves, flowers and pods being free from anthocyanin.--

--22. The plant of claim 16 wherein the apex of said leaves are acuminate and the base of said leaves is obtuse.--

--23. The plant of claim 22, wherein the average height of said plant when mature is about 34.9 cm.--

--24. The plant of claim 22, wherein said plant has lodging resistance through maturity and withstands wind and other climatic conditions.--

--25. The plant of claim 22, wherein said plant establishes a long deep-growing wood-like taproot, a plurality of wood-like lateral roots, and a plurality of wood-like feeder roots.--

--26. The plant claim 25, wherein said taproot is larger than at least one of said lateral roots, and at least one of said lateral roots is larger than said feeder roots.--

--27. The plant of claim 25, wherein said taproot averages 1.0 cm +/- in caliper size.--

2001-07-16



RECEIVED

JUL 19 2001

TECH CENTER 1600/2900

--28. A pod of a *Phaseolus vulgaris* field bean plant having, at onset, a solid green color pattern, wherein said color is about 5 GY 6/6 in the *Munsell Book of Color* when viewed in natural light.--

--29. The pod of claim 28, wherein said pod has a pear-shaped cross section.--

--30. The pod of claim 29, wherein the curvature of said pod is straight and the orientation of the beak of said pod is straight.--

--31. The pod of claim 30, wherein said pod has slight constrictions.--

--32. A pod of a *Phaseolus vulgaris* field bean plant having, at maturity, a solid tan color pattern, wherein said color is about 5 Y 8.5/6 in the *Munsell Book of Color* when viewed in natural light.--

--33. The pod of claim 32, wherein said pod has a pear-shaped cross section.--

--34. The pod of claim 32, wherein said pod is slightly curved and the orientation of the beak of said pod is variable.--

--35. The pod of claim 32, wherein said pod has slight constrictions.--

--36. The pod of claim 32, wherein the average beak length of said pod is 1.2 cm.--

--37. The pod of claim 32, wherein said pod comprises seeds and the number of said seeds per pod is approximately 3.1.--

--38. A *Phaseolus vulgaris* field bean plant, said plant comprising a wood-like stalk, at least one pod, and a plurality of wrinkled, dull, ovate-shaped leaves.--

al
1600-2900



RECEIVED

JUL 19 2001

TECH. CENTER 1600/2900

--39. The plant of claim 38, wherein said pod has, at onset, a solid green color pattern, wherein said color is about 5 GY 6/6 in the *Munsell Book of Color* when viewed in the natural light.--

--40. The plant of claim 39, wherein said pod has a pear-shaped cross section.--

--41. The plant of claim 40, wherein said pod is straight and the orientation of the beak of said pod is straight.--

--42. The plant of claim 41, wherein said pod has slight constrictions.--

--43. The plant of claim 38, wherein said pod has, at maturity, a solid tan color pattern, wherein said color is about 5 Y 8.5/6 in the *Munsell Book of Color* when viewed in natural light.--

--44. The plant of claim 43, wherein said pod has a pear-shaped cross section.--

--45. The plant of claim 43, wherein said pod is slightly curved and the orientation of the beak of said pod is variable.--

--46. The plant of claim 43, wherein said pod has slight constrictions.--

--47. The plant of claim 43, wherein the average beak length of said pod is 1.2 cm.--

--48. The plant of claim 43, wherein said pod comprises seeds and the number of said seeds per pod is approximately 3.1.--

--49. A method of harvesting a *Phaseolus vulgaris* field bean plant, said method comprising the following steps:

- a. knifing the plant,
- b. placing the plant into a window;

a1

1000-0000-0000-0000



RECEIVED

JUL 19 2001

TECH CENTER 1600/2900

c. allowing the plant to dry.--

--50. The method of claim 49 wherein said drying step is continued for approximately 5 to 8 days.--

Sub 7
C27
--51. A seed from a field bean variety of *Phaseolis vulgaris* comprising a seed coat and a hilar ring wherein the seed coat color is about 7.5 Y 8.5/4 to about 7.5 Y 8.5/6 in the *Munsell Book of Color* when viewed in natural light.--

--52. The seed of claim 51 wherein the hilar ring color is from about 2.5 Y 9/4 to about 2.5 Y 9/6 in the *Munsell Book of Color* when viewed in natural light.--

at
Sub 3
SC3
--53. A seed from a field bean variety of *Phaseolis vulgaris* comprising a seed coat and a hilar ring wherein the hilar ring color is from about 2.5 Y 9/4 to about 2.5 Y 9/6 in the *Munsell Book of Color* when viewed in natural light.--

--54. The seed of claim 51, wherein said seed, when eaten, has a smooth texture and pleasing taste.--

--55. The seed of claim 51, wherein said seed is able to take on a large volume of water when soaked prior to cooking.--

--56. The seed of claim 51, wherein germination occurs in an environment free of light.--

--57. The seed of claim 51, wherein said seed is grown in a pod and the shape of said seed taken from the middle of said pod is cuboid.--

--58. The seed of claim 51, wherein the dry seed weight is about 43 grams per 100 seeds (adjust to 12 percent moisture).--